



Contents lists available at ScienceDirect

International Journal of Infectious Diseases

journal homepage: [www.elsevier.com/locate/ijid](http://www.elsevier.com/locate/ijid)

# Impact of the International Nosocomial Infection Control Consortium (INICC) multidimensional hand hygiene approach in three cities of Colombia<sup>☆</sup>



Nayide Barahona-Guzmán<sup>a</sup>, María Eugenia Rodríguez-Calderón<sup>b</sup>, Victor D. Rosenthal<sup>c,\*</sup>, Narda Olarte<sup>d</sup>, Wilmer Villamil-Gómez<sup>e,f</sup>, Catherine Rojas<sup>g</sup>, Marena Rodríguez-Ferrer<sup>a</sup>, Guillermo Sarmiento-Villa<sup>a</sup>, Alfredo Lagares-Guzmán<sup>a</sup>, Alberto Valderrama<sup>d</sup>, Antonio Menco<sup>e</sup>, Patrick Arrieta<sup>e,f</sup>, Luis Enrique Dajud-Cassas<sup>f</sup>, Mariela Mendoza<sup>f</sup>, Alejandra Sabogal<sup>g</sup>, Yulieth Carvajal<sup>g</sup>, Edwin Silva<sup>g</sup>

<sup>a</sup> Universidad Simón Bolívar, Barranquilla, Colombia

<sup>b</sup> Hospital La Victoria, Bogotá, Colombia

<sup>c</sup> International Nosocomial Infection Control Consortium (INICC), Corrientes Ave 4580, Floor 12, Apt D, Buenos Aires, ZIP 1195, Argentina

<sup>d</sup> Hospital El Tunal ESE, Bogotá, Colombia

<sup>e</sup> Clínica Santa María, Sucre, Colombia

<sup>f</sup> Clínica de la Sabana, Sucre, Colombia

<sup>g</sup> Centro Policlínico del Olaya, Bogotá, Colombia

## ARTICLE INFO

### Article history:

Received 9 May 2013

Received in revised form 28 October 2013

Accepted 28 October 2013

**Corresponding Editor:** Eskild Petersen, Aarhus, Denmark

### Keywords:

Hand hygiene

Intensive care unit

Bundle

International Nosocomial Infection Control Consortium

Multidimensional approach

Developing countries

## SUMMARY

**Objective:** To assess the feasibility and effectiveness of the International Nosocomial Infection Control Consortium (INICC) multidimensional hand hygiene (HH) approach in Colombia, and analyze predictors of poor HH compliance.

**Methods:** An observational, prospective, interventional, before-and-after study was conducted from May 2003 through September 2010 in 10 intensive care units (ICUs) of six hospitals in three cities. The study was divided into two periods: a baseline and a follow-up period. Observations for HH compliance were done in each ICU during randomly selected 30-min periods. The multidimensional HH approach included: (1) administrative support, (2) supplies availability, (3) education and training, (4) reminders in the workplace, (5) process surveillance, and (6) performance feedback.

**Results:** A total of 13 187 opportunities for HH were observed. Overall HH compliance increased from 50% to 77% (relative risk 1.55, 95% confidence interval 1.43–1.68;  $p = 0.0001$ ). Multivariate and univariate analyses showed that several variables were significantly associated with poor HH compliance: males vs. females (67% vs. 77%;  $p = 0.0001$ ), physicians vs. nurses (59% vs. 78%;  $p < 0.0001$ ), and adult vs. pediatric ICUs (76% vs. 42%;  $p < 0.001$ ), among others.

**Conclusions:** Adherence to HH was increased by 55% with the INICC approach. Programs targeted at improving HH in variables found to be predictors of poor compliance should be implemented.

© 2013 The Authors. Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. All rights reserved.

## 1. Introduction

Back in the nineteenth century, when Semmelweis studied the relationship between improved hand antisepsis and reduced mortality from puerperal sepsis,<sup>1</sup> it was shown that appropriate hand hygiene (HH) before patient contact was a fundamental tool for the prevention of infection. Since then, several studies have reported that improved HH practice is associated with a reduction

<sup>☆</sup> This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-No Derivative Works License, which permits non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

\* Corresponding author. Tel./Fax: +54 11 4861 5826.

E-mail address: [victor\\_rosenthal@inicc.org](mailto:victor_rosenthal@inicc.org) (V.D. Rosenthal).

URL: <http://www.inicc.org>

in the rates of healthcare-associated infection (HAI) and antimicrobial resistance.<sup>2–4</sup>

HAIs pose a threat to patient safety, including morbidity and mortality.<sup>5</sup> Traditionally, most studies focusing on HAIs have been conducted in developed countries;<sup>6</sup> in developing countries, this public health problem had not been systematically studied until the International Nosocomial Infection Control Consortium (INICC) began to measure and analyze HAI rates with standardized definitions and methods.<sup>7–11</sup>

HH practice plays a crucial role in preventing cross-transmission of HAIs, and successful interventions to improve HH have been reported in the mainstream literature from developed<sup>12</sup> and developing nations.<sup>3,13,14</sup> Since the 1980s, investigators have analyzed the effectiveness of several interventions to improve HH, such as the use of monitoring and performance feedback as published by Mayer et al. in 1986;<sup>15</sup> the impact of supplies availability as published by Preston et al. in 1981;<sup>16</sup> administrative support as published by Larson et al. in 1997;<sup>17</sup> the effectiveness of education as published by Dubbert et al. in 1990,<sup>18</sup> and by Dorsey et al. in 1996;<sup>19</sup> the use of reminders and posters in the workplace as published by Conly et al. in 1989;<sup>20</sup> and the introduction of alcohol-based hand rub (AHR) as published by Graham in 1990.<sup>21</sup> In a study conducted in the USA in 1997, Larson et al. explicitly referred to a multidimensional approach that considered several interventions.<sup>17</sup> Likewise, Rosenthal et al. have implemented programs in Argentina since 1993, combining administrative support, supplies availability, education and training, process surveillance, and performance feedback, which produced a sustained improvement in HH compliance,<sup>14</sup> with a reduction in HAI rates.<sup>3</sup>

In 2002, HH guidelines were published by the US Centers for Disease Control and Prevention (CDC).<sup>22</sup> With a view to promoting HH from a global angle, in 2005 the World Health Organization (WHO) launched the program “Clean Care is Safer Care”,<sup>23</sup> and in 2009 the WHO published its guidelines presenting a compilation of previously published data, and a new formulation for AHR products, among several other recommendations.<sup>4</sup>

There are no previous publications showing HH compliance in the hospitals of Colombia. The aim of this study was to establish the baseline HH compliance rate of healthcare workers (HCWs) before patient contact in 10 intensive care units (ICUs) of six hospitals in three cities of Colombia, to analyze risk factors for poor adherence, and to evaluate the impact of implementing the INICC multidimensional HH approach (IMHHA). The IMHHA includes the following elements: (1) administrative support, (2) supplies availability, (3) education and training, (4) reminders in the workplace, (5) process surveillance, and (6) performance feedback.

## 2. Methods

### 2.1. The INICC

The INICC is an open, non-profit, HAI surveillance multicenter network that applies methods based on the US CDC/National Healthcare Safety Network (NHSN).<sup>24</sup> The INICC was established to measure and control HAIs in hospitals worldwide through the analysis of standardized data collected on a voluntary basis by its member hospitals and fosters the use of evidence-based preventive measures. Since its international inception in 2002, the INICC has increasingly gained new members and now comprises nearly 1000 hospitals in 200 cities of 50 countries in Latin America, Asia, Africa, the Middle East, and Europe; it has become the only source of aggregate standardized international data on the epidemiology of HAIs internationally.<sup>11</sup>

### 2.2. Study setting

This study was conducted in 10 ICUs of six hospitals in three cities of Colombia, which were successively incorporated into the study over a period of 8 years.

Each hospital has an infection control team (ICT) with at least one infection control practitioner (ICP) and one physician, but the number of members varies depending on the ICU. The ICT member in charge of process surveillance at each hospital has at least 2 years of experience in monitoring HAI rates and infection control practices.

Professional categories of HCWs included nurses, physicians, and ancillary staff (including paramedical technicians, nurse aides, laboratory team members, radiology team members, physiotherapists, patient care technicians, paramedical personnel, and patient lift teams).

The study protocol was approved by the institutional review boards at each hospital.

### 2.3. Study design

An observational, prospective, cohort, interventional, before-and-after multicenter study was conducted from May 2003 through September 2010. The study was divided into two periods: a baseline and a follow-up period. The baseline period for HH compliance included episodes documented at each hospital during their first 3 months of participation, and the follow-up period included episodes following the fourth month of participation. Each ICU started to participate in the study at different times, and therefore have different lengths of follow-up; however the length of the baseline period was exactly the same (3 months) for all ICUs. For comparison of the compliance rates, the ICUs were aligned independently of the date at which they started to participate in the study.

### 2.4. INICC multidimensional HH approach (IMHHA)

The IMHHA is implemented at each hospital from the beginning of their participation in the INICC. This includes the following six components: (1) administrative support, (2) supplies availability, (3) education and training, (4) reminders in the workplace, (5) process surveillance, and (6) performance feedback. Although the components are presented individually, they are interactive elements that must occur together for the effective implementation of any ‘multidimensional’ approach.

#### 2.4.1. Administrative support

Hospital administrators at the participating hospitals agreed and committed to the study, attended infection control meetings to discuss study findings, and allocated supplies of HH products.

#### 2.4.2. Supplies availability

During the study period, AHR bottles were available at the ICU entrances, nursing stations, and near the site of patient care (individual patient room entrances, at bedside tables, and/or at the foot of the patient’s bed). Sinks with a water supply, soap, and paper towels were available at the ICU entrances, nursing stations, and common areas of the ICUs.

#### 2.4.3. Education and training

At the study ICUs, the ICT members provided 30-min education sessions to HCWs of each work shift, at the beginning of the study period and at regular times periodically during the follow-up period (every month, every 2 months, and every 6 months, depending on the ICU). Education included information about indications for HH and the correct procedures and techniques for HH.

**Table 1**  
Characteristics of the participating hospitals and intensive care units

|                  | n (%)    | HH baseline observations, n | HH intervention observations, n | HH overall observations, n |
|------------------|----------|-----------------------------|---------------------------------|----------------------------|
| Type of ICU      |          |                             |                                 |                            |
| Coronary         | 1 (10%)  | 132                         | 1383                            | 1515                       |
| Medical surgical | 5 (50%)  | 553                         | 9148                            | 9701                       |
| Newborn          | 2 (20%)  | 381                         | 1142                            | 1523                       |
| Neurosurgical    | 1 (10%)  | 127                         | 76                              | 203                        |
| Pediatric        | 1 (10%)  | 101                         | 144                             | 245                        |
| All ICUs         | 10       | 1294                        | 11 893                          | 13 187                     |
| Type of hospital |          |                             |                                 |                            |
| Private          | 4 (67%)  | 752                         | 10 772                          | 11 524                     |
| Public           | 2 (33%)  | 542                         | 1121                            | 1663                       |
| All hospitals    | 6 (100%) | 1294                        | 11 893                          | 13 187                     |

HH, hand hygiene; ICU, intensive care unit.

#### 2.4.4. Reminders in the workplace

Poster reminders were displayed all around the hospital setting (i.e., hospital entrance, corridors, ICT office, ICU entrances, nursing stations, beside each sink, and beside each AHR bottle). They included simple instructions on HH performance, in line with the content of the education and training program.

#### 2.4.5. Process surveillance

Process surveillance of HH practices consisted of the registration of potential opportunities for HH,<sup>4</sup> and the actual number of HH episodes, either with water and soap or AHR. HCW HH practice was monitored directly by an observer, a member of the ICT, following a standardized protocol and completion of HH process surveillance INICC forms.<sup>7</sup> Observations were conducted unobtrusively at specific time periods distributed across three work shifts (morning, afternoon, and evening). HCWs were not aware of the schedule of the monitoring period. The monitoring included HH compliance before patient contact and before an aseptic task, because at the time we started the study in May 2003, the “Five Moments for Hand Hygiene” proposed by the WHO had not been published; they were published in 2009. Potential confounders of HH included sex, type of ICU, professional category, work shift, and type of contact.

#### 2.4.6. Performance feedback

Every month, the INICC Headquarters team prepares and sends a final month-by-month report on compliance with HH to each participating ICU. These charts contain a running tally of HH compliance by HCWs of the ICUs, and compliance comparing several variables, such as sex, HCW professional status, ICU type, contact type, and work shift. These charts were reviewed at

monthly ICT meetings and also posted in the ICUs to give performance feedback to the HCWs of the participating ICUs.<sup>7</sup> The performance feedback process started in the third month of participation in this approach.<sup>7</sup>

#### 2.5. Training of the infection control team for process surveillance

The ICT investigators were self-trained with a procedure manual sent from the INICC Headquarters in Buenos Aires, specifying how to carry out the HH process surveillance and how to fill in the INICC forms.<sup>7</sup> ICT members had continuous telephone, e-mail, and webinar access to a support team at the INICC Headquarters.


#### 2.6. Data collection and processing

Completed INICC process surveillance forms for HH were sent monthly by the ICT members from each participating ICU to the INICC Headquarters. The team at the INICC Headquarters uploaded the data into a database, performed an analysis, and sent a report on HH compliance to the ICT members of each participating ICU, showing HH compliance by month, by sex, by HCW profession, by ICU, by work shift, and by type of contact.<sup>7</sup>

#### 2.7. Statistical methods

##### 2.7.1. Univariate analysis of variables associated with poor HH and of the impact of the HH approach

The aggregated independent variables (sex and profession of HCWs, type of ICU, type of contact, etc.) of all observed HH opportunities and HH compliance during the whole study, and a



**Figure 1.** Types of product used in hand hygiene over the years of participation.

**Table 2**  
Distribution of hand hygiene compliance by type of intensive care unit

| ICU type         | Number of ICUs | Baseline period<br>(HH compliance/HH observations) | Intervention period<br>(HH compliance/HH observations) | RR (95% CI)      | p-Value |
|------------------|----------------|--|--|------------------|---------|
| Coronary         | 1              | 22.7% (30/132)                                     | 25.5% (352/1383)                                       | 1.12 (0.77–1.63) | 0.5514  |
| Medical surgical | 5              | 62.0% (343/553)                                    | 85.7% (7839/9148)                                      | 1.38 (1.24–1.54) | 0.0001  |
| Newborn          | 1              | 57.2% (218/381)                                    | 79.2% (904/1142)                                       | 1.38 (1.19–1.60) | 0.0001  |
| Neurosurgical    | 1              | 29.1% (37/127)                                     | 44.7% (34/76)  | 1.54 (0.96–2.45) | 0.0688  |
| Pediatric        | 1              | 17.8% (18/101)                                     | 59.7% (86/144)   | 3.35 (2.02–5.57) | 0.0001  |
| All              | 9              | 49.9% (646/1294)                                   | 77.5% (9215/11 893)                                    | 1.55 (1.43–1.68) | 0.0001  |

ICU, intensive care unit; HH, hand hygiene; RR, relative risk; CI, confidence interval.

comparison of HH compliance during the baseline period and during the follow-up period, were compared using Fisher's exact test for dichotomous variables and the unmatched Student's *t*-test for continuous variables. Relative risk (RR) ratios were calculated for comparisons of analyzed variables associated with HH using EPI Info v. 6. Confidence intervals (95% CI) were calculated using VCStat (Castiglia). *p*-Values of <0.05 by two-tailed tests were considered significant.

### 2.7.2. Multivariate analysis of variables associated with poor HH

The aggregated independent variables described for all observed HH opportunities and HH compliance during the whole study were compared using logistic regression for dichotomous and continuous variables. Odds ratios (OR) with 95% CI were calculated for comparisons of analyzed variables associated with HH using PASW Statistics 18 (SPSS Inc.). *p*-Values of <0.05 by two-tailed tests were considered significant.

### 2.7.3. Multivariate analysis of the impact of the INICC multidimensional HH approach (IMHHA)

HH opportunities and HH compliance during baseline and during follow-up were explored for changes in HH compliance rates following an ICU joining the INICC. We present the results of a logistic regression model to consider the change in HH compliance in INICC participating ICUs over time since the beginning of the HH surveillance. ORs are presented, comparing each time period since the start of the surveillance with the baseline of 3 months. Because of the different lengths of follow-up at each ICU, for each time period only ICUs with follow-up during that time period were included in the baseline period used for calculating the OR of HH compliance for that period.

**Table 3**  
Hand hygiene compliance by type of variable: univariate analysis

| Data             | Variable        | % (number of HH/number opportunities) | Comparison | RR   | 95% CI    | p-Value |
|------------------|-----------------|---------------------------------------|------------|------|-----------|---------|
| Sex              | Female          | 77% (8024/10 446)                     | F vs. M    | 0.87 | 0.83–0.92 | 0.0001  |
|                  | Male            | 67% (1837/2741)                       |            |      |           |         |
| HCW              | Nurses          | 78% (5198/6656)                       | Ns vs. Ph  | 0.76 | 0.71–0.81 | 0.0001  |
|                  | Physicians      | 59% (1127/1908)                       | Ns vs. AS  | 0.98 | 0.94–1.02 | 0.3399  |
|                  | Ancillary staff | 76% (3536/4623)                       | Ph vs. AS  | 0.77 | 0.72–0.83 | 0.0001  |
| Procedure        | Non-invasive    | 71% (6360/8956)                       | NI vs. I   | 0.84 | 0.88–0.87 | 0.0001  |
|                  | Invasive        | 85% (3337/3939)                       |            |      |           |         |
| ICU              | Adult           | 76% (8635/11 419)                     | Ad vs. Pe  | 0.56 | 0.46–0.68 | 0.0001  |
|                  | Newborn         | 74% (1112/1523)                       | Ad vs. Nb  | 0.97 | 0.92–1.04 | 0.4105  |
|                  | Pediatric       | 42% (104/245)                         | Nb vs. Pe  | 0.58 | 0.47–0.70 | 0.0001  |
| Work shift       | Morning         | 78% (5299/6816)                       | M vs. A    | 0.94 | 0.89–0.98 | 0.0036  |
|                  | Afternoon       | 73% (2951/4058)                       | M vs. N    | 0.90 | 0.85–0.95 | 0.0001  |
|                  | Night           | 70% (1611/2313)                       | A vs. N    | 0.96 | 0.90–1.02 | 0.1636  |
| Type of hospital | Public          | 78% (1299/1663)                       | Pu vs. Pr  | 0.95 | 0.90–1.01 | 0.0926  |
|                  | Private         | 74% (8562/11 524)                     |            |      |           |         |

HH, hand hygiene; RR, relative risk; CI, confidence interval; HCW, healthcare worker; ICU, intensive care unit.

## 3. Results

From May 2003 to September 2010, we recorded a total 13 187 opportunities for HH before patient contact and before an aseptic task. Characteristics of the hospitals and ICUs participating in the study from May 2003 to September 2010 are shown in Table 1.

The distribution of use of HH products over the study period is shown in Figure 1.

### 3.1. Predictors of poor HH compliance

We observed 11 524 procedures in private hospitals and 1663 in public hospitals; 10 446 procedures in females and 2741 in males; 6656 in nurses, 1908 in physicians, and 4623 in ancillary staff; 8956 were prior to non-invasive patient contacts and 3939 prior to invasive procedures; 11 419 in adult ICUs, 245 in pediatric ICUs (PICUs), and 1523 in neonatal ICUs (NICUs); 6816 during the morning, 4058 during the afternoon, and 2313 during the night shift.

Table 2 shows the distribution of HH compliance among the different ICU types in the baseline and intervention periods.

Tables 3 and 4 show HH compliance according to each variable (type of hospital, sex, profession of HCW, type of procedure, type of unit, and work shift), and the association with poor HH, analyzed with univariate and multivariate statistical methods.

### 3.2. Components of the INICC multidimensional HH approach (IMHHA)

During the follow-up period, the six components of the IMHHA were applied simultaneously: 100% counted on administrative

**Table 4**  
Hand hygiene compliance by type of variable: logistic regression, multivariate analysis

| Variable                                | Adjusted OR (95% CI) | p-Value |
|---|----------------------|---------|
| Type of hospital (baseline: Public)     | 1                    |         |
| Private                                 | 0.65 (0.54–0.80)     | 0.001   |
| Sex (baseline: Female)                  | 1                    |         |
| Male                                    | 0.88 (0.79–0.98)     | 0.024   |
| Type of professional (baseline: Nurses) | 1                    |         |
| Physicians                              | 0.41 (0.36–0.47)     | 0.001   |
| Ancillary staff                         | 0.92 (0.84–1.01)     | 0.092   |
| Type of contact (baseline: Invasive)    | 1                    |         |
| Non-invasive                            | 0.42 (0.38–0.47)     | 0.001   |
| Type of ICU (baseline: Adult)           | 1                    |         |
| Adult ICU                               | 0.56 (0.46–0.69)     | 0.001   |
| Pediatric ICU                           | 0.28 (0.22–0.37)     | 0.001   |
| Work shift (baseline: Morning)          | 1                    |         |
| Afternoon                               | 0.77 (0.71–0.85)     | 0.001   |
| Night                                   | 0.70 (0.63–0.78)     | 0.001   |

OR, odds ratio; CI, confidence interval; ICU, intensive care unit.

support and available supplies for HH and AHR; 100% educated HCWs (14.3% of them every month, 57.1% every 2 months, 14.3% every 3 months, and 14.3% every 6 months); 100% posted reminders (100% of them at ICU entrances, 100% in common ICU areas, 36.4% beside each bed); process surveillance was conducted by 100%; 100% provided performance feedback (14.3% of them every month, 28.6% every 2 months, 28.6% every 3 months).

### 3.3. Impact of the INICC multidimensional HH approach (IMHHA) on HH compliance

The baseline period of the INICC ICUs was 3 months and their average follow-up period was 42.0 months (range 4–84 months). The results of a logistic regression model to consider the change in HH compliance in INICC participating ICUs over the whole study period are presented in Table 5 and Figure 2.

## 4. Discussion

This is the first study to show an improvement in HH compliance in Colombia due to the implementation of the IMHHA. We analyzed the impact of the IMHHA in a diverse ICU population in three cities of Colombia, showing that the six measures of the IMHHA implemented in each ICU were followed by a 55% increase in the adherence to proper HH practices.

Baseline HH compliance (50%) of HCWs in the INICC ICUs was similar to that shown in previous studies, although HH compliance rates have ranged from 9% to 75% in different studies.<sup>4</sup>

The results of the multivariate analysis showed that there was higher compliance in females, which has also been identified in individuals unrelated to healthcare: Guinan et al. showed higher HH compliance by female students.<sup>25</sup> In consonance with this, compliance was also higher among nurses than physicians, as also shown in a study by Rosenthal et al. in 2005.<sup>3</sup> We also showed the highest HH compliance was in the adult ICUs. Our findings show that the type of contact influenced HH performance: non-invasive

contact was associated with lower compliance than invasive contact. This finding is in agreement with Lipsett and Swoboda, who suggested that low-risk situations were predictors for lower compliance.<sup>26</sup>

Our approach included administrative support; in 2003 Rosenthal et al. showed that higher HH adherence was associated with administrative support.<sup>14</sup> It also included supplies availability, which was shown to improve HH compliance.<sup>14</sup> The IMHHA also included education and training, which are other basic independent interventions identified as fostering adequate HH performance. As shown in 1990 by Dubbert et al.,<sup>18</sup> educational interventions with routine classes improved HH compliance by 97% over 4 weeks. Likewise, but within the context of resource-limited countries, Rosenthal et al.<sup>14</sup> showed HCW education improved HH adherence and that compliance increased further if performance feedback was also implemented. We also included reminders in the workplace. In 1989, Conly et al.<sup>20</sup> showed the importance of reminders to raise HCW awareness of the relationship between correct HH performance and the reduction of HAI.

We measured 13 187 opportunities for HH. On a monthly basis, the ICT team provided performance feedback to the HCWs of each ICU. This is a highly motivating aspect of the IMHHA for HCWs. Knowing the outcome of their efforts reflected by the measurement of their practices and the incidence of HAIs can be a most rewarding or awareness-raising factor and ensures the effectiveness of the IMHHA.<sup>27,28</sup> Since 1998 in Argentina,<sup>3,14</sup> and 2002 internationally,<sup>7–11</sup> the INICC has introduced outcome and process surveillance and feedback on outcomes and performance, combined with training and education, as a means to improve quality in healthcare to a new level.<sup>3,14</sup>

Through the last decade, the INICC has undertaken a global effort in America, Asia, Africa, the Middle East, and Europe to respond to the burden of HAIs, and has achieved successful results by increasing HH compliance, improving compliance with other infection control interventions, as described in several INICC publications, and consequently, reducing the rates of HAI and mortality. Since 2002, in adult ICUs in 15 countries, the INICC has reduced the rate of central line-associated bloodstream infection (CLAB) by 54%,<sup>29</sup> of catheter-associated urinary tract infection (CAUTI) by 37%,<sup>30</sup> of ventilator-associated pneumonia (VAP) by 56%,<sup>31</sup> and of mortality by 58%.<sup>29</sup> In PICUs in five countries, the INICC have reduced the rate of CLAB by 52%,<sup>32</sup> of CAUTI by 57%,<sup>33</sup> of VAP by 31%,<sup>34</sup> and of mortality by 31%.<sup>32</sup> In NICUs in 10 countries, the INICC has reduced the rate of VAP by 33%.<sup>35</sup>

This study has several limitations. First, we did not measure the “Five Moments for Hand Hygiene” as recently described by the WHO in 2009. This is because the INICC started the IMHHA in 1998 in Argentina,<sup>3,14</sup> which was launched internationally in 2002,<sup>7</sup> i.e., several years before the WHO published its recommendations in 2009. However, since 2009, the INICC has included the WHO recommendations in its process surveillance forms and manuals.<sup>4</sup> Additionally, it should be noted that this study applied an observational, before-and-after methodology, which implies less strength of evidence than other study designs. A Hawthorne effect

**Table 5**  
Hand hygiene improvement by year of participation

| Years since joining INICC | HH observations | Number of ICUs included | Number of hospitals included | HH % (95% CI)     | Adjusted OR (95% CI) | p-Value |
|---------------------------|-----------------|-------------------------|------------------------------|-------------------|----------------------|---------|
| First 3 months (baseline) | 1294            | 10                      | 6                            | 49.9% (0.47–0.52) | 1.0                  |         |
| Months 4–6                | 823             | 10                      | 6                            | 57.8% (0.54–0.61) | 1.04 (0.84–1.31)     | 0.694   |
| Months 7–9                | 761             | 8                       | 5                            | 68.6% (0.65–0.72) | 2.45 (1.9–3.2)       | 0.001   |
| Months 10–12              | 767             | 8                       | 5                            | 67.9% (0.65–0.71) | 2.83 (2.2–3.7)       | 0.001   |
| Second and third year     | 9542            | 5                       | 4                            | 80.7% (0.80–0.81) | 4.70 (3.6–6.2)       | 0.0001  |

INICC, International Nosocomial Infection Control Consortium; HH, hand hygiene; ICU, intensive care unit; CI, confidence interval; OR, odds ratio.




Figure 2. Hand hygiene improvement by year of participation.

is typical in direct observations of adherence. In addition, direct observations represent only a sample of all opportunities and there are inherent weaknesses including assuring inter-observer reliability. Finally, capturing the quality of the HH technique is highly complex, and we were not able to include many details in this study, such as information regarding HAI and mortality rates, since there are several INICC publications that have focused on these topics in relation to HH.

In conclusion, we found that the IMHHA improved HH compliance in 10 ICUs of six hospitals in three cities of Colombia. It is the primary objective of the INICC to foster infection control practices by freely facilitating elemental and inexpensive resource tools to tackle this problem effectively and systematically, leading to greater and steady adherence to infection control programs and guidelines, such as HH compliance, and to the correlated reduction in HAIs and their consequences, such as extra mortality and costs.

## Acknowledgments

The authors thank the many healthcare professionals at each member hospital who assisted in the conducting of surveillance in their hospital, including the surveillance nurses, clinical microbiology laboratory personnel, and the physicians and nurses providing care for the patients during the study; without their cooperation and generous assistance this INICC would not be possible. We also thank Mariano Vilar and Débora López Burgardt, who work at INICC Headquarters in Buenos Aires, for their hard work and commitment to achieve INICC goals; the INICC country coordinators and INICC Advisory Board, who have so generously supported this unique international infection control network.

**Funding:** The funding for the activities carried out at INICC Headquarters was provided by the corresponding author, Victor D. Rosenthal, and the Foundation to Fight against Nosocomial Infections.

**Ethical approval:** Every hospital's institutional review board agreed to the study protocol. Patient confidentiality was protected by codifying the recorded information, making it only identifiable to the infection control team.

**Conflict of interest:** All authors report no conflicts of interest related to this article.

## References

1. Raju TN. Ignac Semmelweis and the etiology of fetal and neonatal sepsis. *J Perinatol* 1999;**19**:307–10.

2. Simmons B, Bryant J, Neiman K, Spencer L, Arheart K. The role of handwashing in prevention of endemic intensive care unit infections. *Infect Control Hosp Epidemiol* 1990;**11**:589–94.
3. Rosenthal VD, Guzman S, Safdar N. Reduction in nosocomial infection with improved hand hygiene in intensive care units of a tertiary care hospital in Argentina. *Am J Infect Control* 2005;**33**:392–7.
4. Pittet D, Allegranzi B, Boyce J. The World Health Organization Guidelines on Hand Hygiene in Health Care and their consensus recommendations. *Infect Control Hosp Epidemiol* 2009;**30**:611–22.
5. Jarvis WR. Selected aspects of the socioeconomic impact of nosocomial infections: morbidity, mortality, cost, and prevention. *Infect Control Hosp Epidemiol* 1996;**17**:552–7.
6. Safdar N, Crnich CJ, Maki DG. Nosocomial infections in the intensive care unit associated with invasive medical devices. *Curr Infect Dis Rep* 2001;**3**:487–95.
7. Rosenthal VD, Maki DG, Graves N. The International Nosocomial Infection Control Consortium (INICC): goals and objectives, description of surveillance methods, and operational activities. *Am J Infect Control* 2008;**36**:e1–2.
8. Rosenthal VD, Maki DG, Salomao R, Moreno CA, Mehta Y, Higuera F, et al. Device-associated nosocomial infections in 55 intensive care units of 8 developing countries. *Ann Intern Med* 2006;**145**:582–91.
9. Rosenthal VD, Maki DG, Mehta A, Alvarez-Moreno C, Leblebicioglu H, Higuera F, et al. International Nosocomial Infection Control Consortium report, data summary for 2002–2007, issued January 2008. *Am J Infect Control* 2008;**36**:627–37.
10. Rosenthal VD, Maki DG, Jamulitrat S, Medeiros EA, Todi SK, Gomez DY, et al. International Nosocomial Infection Control Consortium (INICC) report, data summary for 2003–2008, issued June 2009. *Am J Infect Control* 2010;**38**:95–104.e2.
11. Rosenthal VD, Bijie H, Maki DG, Mehta Y, Apisarnthanarak A, Medeiros EA, et al. International Nosocomial Infection Control Consortium (INICC) report, data summary of 36 countries, for 2004–2009. *Am J Infect Control* 2012;**40**:396–407.
12. Lam BC, Lee J, Lau YL. Hand hygiene practices in a neonatal intensive care unit: a multimodal intervention and impact on nosocomial infection. *Pediatrics* 2004;**114**:e565–71.
13. Allegranzi B, Sax H, Bengaly L, Richet H, Minta DK, Chrait MN, et al. Successful implementation of the World Health Organization hand hygiene improvement strategy in a referral hospital in Mali, Africa. *Infect Control Hosp Epidemiol* 2010;**31**:133–41.
14. Rosenthal VD, McCormick RD, Guzman S, Villamayor C, Orellano PW. Effect of education and performance feedback on handwashing: the benefit of administrative support in Argentinean hospitals. *Am J Infect Control* 2003;**31**:85–92.
15. Mayer JA, Dubbert PM, Miller M, Burkett PA, Chapman SW. Increasing handwashing in an intensive care unit. *Infect Control* 1986;**7**:259–62.
16. Preston GA, Larson EL, Stamm WE. The effect of private isolation rooms on patient care practices, colonization and infection in an intensive care unit. *Am J Med* 1981;**70**:641–5.
17. Larson EL, Bryan JL, Adler LM, Blane C. A multifaceted approach to changing handwashing behavior. *Am J Infect Control* 1997;**25**:3–10.
18. Dubbert PM, Dolce J, Richter W, Miller M, Chapman SW. Increasing ICU staff handwashing: effects of education and group feedback. *Infect Control Hosp Epidemiol* 1990;**11**:191–3.
19. Dorsey ST, Cydulka RK, Emerman CL. Is handwashing teachable? Failure to improve handwashing behavior in an urban emergency department. *Acad Emerg Med* 1996;**3**:360–5.
20. Conly JM, Hill S, Ross J, Lertzman J, Louie TJ. Handwashing practices in an intensive care unit: the effects of an educational program and its relationship to infection rates. *Am J Infect Control* 1989;**17**:330–9.
21. Graham M. Frequency and duration of handwashing in an intensive care unit. *Am J Infect Control* 1990;**18**:77–81.

22. Boyce JM, Pittet D. Guideline for hand hygiene in health-care settings. Recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. Society for Healthcare Epidemiology of America/Association for Professionals in Infection Control/Infectious Diseases Society of America. *MMWR Recomm Rep* 2002;**51**:1–45. quiz CE1–4.
23. WHO launches global patient safety challenge; issues guidelines on hand hygiene in health care. *Indian J Med Sci* 2005;**59**:461–3.
24. Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control* 2008;**36**:309–32.
25. Guinan ME, McGuckin-Guinan M, Severeid A. Who washes hands after using the bathroom? *Am J Infect Control* 1997;**25**:424–5.
26. Lipsett PA, Swoboda SM. Handwashing compliance depends on professional status. *Surg Infect (Larchmt)* 2001;**2**:241–5.
27. Bischoff WE, Reynolds TM, Sessler CN, Edmond MB, Wenzel RP. Handwashing compliance by health care workers: the impact of introducing an accessible, alcohol-based hand antiseptic. *Arch Intern Med* 2000;**160**:1017–21.
28. Deming W. Out of the crisis. Boston: MIT Press; 1986.
29. Rosenthal VD, Maki DG, Rodrigues C, Alvarez-Moreno C, Leblebicioglu H, Sobreyra-Oropeza M, et al. Impact of International Nosocomial Infection Control Consortium (INICC) strategy on central line-associated bloodstream infection rates in the intensive care units of 15 developing countries. *Infect Control Hosp Epidemiol* 2010;**31**:1264–72.
30. Rosenthal VD, Todi SK, Alvarez-Moreno C, Pawar M, Karlekar A, Zeggwagh AA, et al. Impact of a multidimensional infection control strategy on catheter-associated urinary tract infection rates in the adult intensive care units of 15 developing countries: findings of the International Nosocomial Infection Control Consortium (INICC). *Infection* 2012;**40**:517–26.
31. Rosenthal VD, Rodrigues C, Alvarez-Moreno C, Madani N, Mitrev Z, Ye G, et al. Effectiveness of a multidimensional approach for prevention of ventilator-associated pneumonia in adult intensive care units from 14 developing countries of four continents: findings of the International Nosocomial Infection Control Consortium. *Crit Care Med* 2012;**40**:3121–8.
32. Rosenthal VD, Ramachandran B, Villamil-Gomez W, Armas-Ruiz A, Navoa-Ng JA, Matta-Cortes L, et al. Impact of a multidimensional infection control strategy on central line-associated bloodstream infection rates in pediatric intensive care units of five developing countries: findings of the International Nosocomial Infection Control Consortium (INICC). *Infection* 2012;**40**:415–23.
33. Rosenthal VD, Ramachandran B, Duenas L, Alvarez-Moreno C, Navoa-Ng JA, Armas-Ruiz A, et al. Findings of the International Nosocomial Infection Control Consortium (INICC), Part I: effectiveness of a multidimensional infection control approach on catheter-associated urinary tract infection rates in pediatric intensive care units of 6 developing countries. *Infect Control Hosp Epidemiol* 2012;**33**:696–703.
34. Rosenthal VD, Alvarez-Moreno C, Villamil-Gomez W, Singh S, Ramachandran B, Navoa-Ng JA, et al. Effectiveness of a multidimensional approach to reduce ventilator-associated pneumonia in pediatric intensive care units of 5 developing countries: International Nosocomial Infection Control Consortium findings. *Am J Infect Control* 2012;**40**:497–501.
35. Rosenthal VD, Rodriguez-Calderon ME, Rodriguez-Ferrer M, Singhal T, Pawar M, Sobreyra-Oropeza M, et al. Findings of the International Nosocomial Infection Control Consortium (INICC), Part II: impact of a multidimensional strategy to reduce ventilator-associated pneumonia in neonatal intensive care units in 10 developing countries. *Infect Control Hosp Epidemiol* 2012;**33**:704–10.